

## Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) from rabbit muscle: Lyophilized

Catalog Number:	1000-E1-GAPDH
Product Specification:	Glyceraldehyde-3-phosphate Dehydrogenase (GAPDH) from rabbit muscle.
Size:	1mg (80 Units)
Type:	Enzyme
Specificity:	One unit will reduce 1 micromole of 3-phosphoglycerate to D-glyceraldehyde 3-phosphate per minute with 3-phosphoglyceric phosphokinase at pH 7.6 at 25 °C.
Western Blot:	The enzyme appears as 36 kDa band when probed with anti-GAPDH antibodies (cat # 1000-M1-GAPDH or 1000-A1-GAPDH)
Immuno-fluorescence:	NA
Immuno-precipitation:	NA
ELISA:	NA
Storage:	-20°C in aliquots. Freeze-thaw cycles must be avoided once the stock aliquot is diluted.
Form:	Lyophilized powder from .

### Product Background:

Glyceraldehyde-3-phosphatedehydrogenase (GAPDH; EC 1.2.1.12) catalyzes an important energy-yielding step in carbohydrate metabolism, the reversible oxidative phosphorylation of glyceraldehyde-3-phosphate in the presence of inorganic phosphate and nicotinamide adenine dinucleotide (NAD). The enzyme is present in such widely separated forms as man, lobster, and *E. coli*. Its rate of evolutionary change is one of the slowest known (1). Recent evidence demonstrates that mammalian GAPDH displays a number of diverse activities unrelated to its glycolytic

function. These include its role in membrane fusion, microtubule bundling, phosphotransferase activity, nuclear RNA export, DNA replication and DNA repair. These new activities may be related to the subcellular localization and oligomeric structure of GAPDH in vivo. Furthermore, other investigations suggest that GAPDH is involved in apoptosis, age-related neurodegenerative disease, prostate cancer and viral pathogenesis. Intriguingly, GAPDH is also a unique target of nitric oxide. Recently GAPDH has been shown to be a common target of a number of compounds with potent antiproliferative activities and thus a potential target for

chemotherapeutic intervention.

Glyceraldehyde-3-phosphate dehydrogenase (EC 1.2.1.12) is also known as GAPDH, G3PDH, OCT1 co-activator in S-phase and p38 component. GAPDH is cytoplasmic, tetrameric (~144 kDa) protein of four identical subunits. Each subunit consists of 333 amino acids (~36 kDa) in mouse and rat and 335 aa in human. The aa seq of the subunit is ~93% identical in human, mouse and rat. The gene encoding GAPDH has been mapped on chromosomes 12 (human) and 6 (mouse). Being a “house-keeping” protein, GAPDH is expressed in almost all the tissues at high levels.

**References:** (1) Michael Sirover, MA (1999) BBA 1432, 159; Xing, C et al (2004) PNAS 101, 5862; Waingeh, VF et al (2004) Biopolymers 73, 533

### Related Products:

1000-A1-GAPDH	Glyceraldehyde-3-phosphate dehydrogenase (GAPDH): Antibody (Rabbit Polyclonal) Aff Pure
1000-D-GAPDH	GAP (Glyceraldehyde-3-Phosphare: Substrate for GAPDH enzyme
1000-E-GAPDH	Glyceraldehyde-3-phosphatedehydrogenase (GAPDH): Enzyme
1000-M1-GAPDH	GAPDH: Antibody (Mouse Monoclonal)
1000-W-GAPDH	Glyceraldehyde-3-phosphatedehydrogenase (GAPDH): Protein (WB +ve Control)

**Disclaimer:** This product is for R&D use only, not for drug, household, or other uses. Dilutions and other information provided are recommendations only; the purchaser must determine the suitability of the product(s) for their particular use. Alamo<sup>®</sup> Laboratories Inc (Alamo<sup>®</sup> Labs) warrants that its product(s) conform to the information contained in this and other Alamo<sup>®</sup> Laboratories Inc publications. If the “Terms and conditions of Sale” are acceptable to you, then you may open the container(s) containing the product and, through such act of opening a container(s), will have shown your acceptance to “Terms and conditions of Sale”. If the terms and conditions are not acceptable to you, then please return the container(s) unopened to Alamo<sup>®</sup> Labs for a complete refund of your payment.